

Short Communication

A Procedure Employing For Redox Titration: Balancing The Redox Chemical Equation In Acidic or Basis Medium

Sneha G. Nair¹, Tushar S. More², Vikram R. Jadhav^{3*}, M. M. Rayate⁴, Bhagyashree A. More⁵

^{2,3}K. K. W. Arts, Science and Commerce College Pimpalgaon (B), Tal - Niphad, Dist - Nashik (India),

^{1,4,5}K. K. Wagh ACS and CS College K-Nagar, Tal - Niphad, Dist - Nashik (India)

Corresponding Author: Vikram R. Jadhav*

ABSTRACT



Graduate and post graduate students find much more difficulty during the concept of the balancing redox chemical equation in acidic or basic medium. Teachers are well aware of the importance of teaching the balancing redox chemical equation in both medium and also difficulty associated with it. In an attempt to improving the understanding of the balancing the redox chemical equation with reduced difficulty, a mathematical approach that provides easier understanding has been developed and successfully employed in graduate and post graduate students. In this study, relation between an oxidising agent (Oxidant) and reducing agent (reductant) can be employed in redox chemical titration, using potassium permanganate (KMnO₄) is an oxidant itself act as the indicator, during the titration of Cr (III) ion, as reducing agent with KMnO₄, the second titration oxidation of H₂O₂, is used as a reducing agent against the KMnO₄. Data obtained from titrations will lead to balancing each redox chemical equation in an acidic or basic medium.

Keywords: Oxidising agent, potassium per magnet, Reducing agent, Redox titration.

INTRODUCTION

A redox chemical equation is an emblematic representation of a chemical reaction. To fulfil the law of conservation of mass these conditions must be balanced. Balanced chemical conditions are basic to take care of issues in the concept of stoichiometry. Fundamentally, balancing redox chemical reaction is a numerical methodology. Most of the redox chemical equations might be balanced by a basic experimentation or assessment method. In any case, for balanced chemical equations are marked as redox reactions, it might appear that there is no basic technique for balancing the redox reactions in acidic or basic medium. There is an enormous number of articles distributed managing an assortment of redox chemical reactions. These range from investigation to logarithmic

technique. In redox conditions, the number of electrons moved from a reducing agent (oxidized species) to an oxidizing agent (reduced species) must be balanced. In general form, a redox chemical equation may be represented as the following way,



m = number of electrons lost by A (Reducing agent)

n = number of electrons gain by B (Oxidising agent)

$m \times \text{moles of reductant} = n \times \text{moles of oxidant.}$


Where, m and n are the stoichiometric coefficients.

$$\frac{m}{n} = \frac{\text{moles of an oxidising agent (oxidant)}}{\text{moles of reducing agent (reductant)}}$$


the above equation indicates that the number of electrons is inversely proportional to moles of respectively agents. To find the ratio of m/n molarity and volume of reducing agent solution used in titration and molarity of MnO_4^- solution are provided.

MATERIALS & METHODS

Part-I: $\text{Cr}^{3+}_{(aq)}/\text{MnO}_4^-_{(aq)}$



The redox chemical reaction of Cr^{3+} with KMnO_4 is presented above by an unbalanced chemical equation in which m and n are the stoichiometric coefficients of reducing and oxidising agent respectively. Here the unbalanced redox reaction may be employed as,



m x moles of reductant $\text{Cr}^{3+} = n$ x moles of oxidant MnO_4^-

$$\frac{m}{n} = \frac{\text{moles of an oxidant}}{\text{moles of reductant}}$$

$$\frac{m}{n} = \frac{\text{moles of an oxidant MnO}_4^-}{\text{moles of reductant Cr}^{3+}}$$

$$w = N \times m \times v$$

$$\frac{m}{n} = \frac{N(\text{MnO}_4^-) \times v(\text{Volume of MnO}_4^-)}{N(\text{H}_2\text{O}_2) \times v(\text{Volume of H}_2\text{O}_2)}$$

$$\frac{m}{n} = \frac{W(\text{MnO}_4^-) \times M(\text{Cr}^{3+})}{M(\text{MnO}_4^-) \times W(\text{Cr}^{3+})}$$


w (Weight of a compound) = N (Normality) x m (molar mass) x v (Volume of a solution)

students performed the titration experiments for the above redox reaction, in which get the following (m/n) ratio in terms of known the Normality of Oxidant (MnO_4^-) and Reductant (Cr^{3+}) and Volume of Cr^{3+} which was taken initially in the titration flask. However, at the


equivalence point the volume of MnO_4^- was considered. The following results was generated

$$\frac{m}{n} = 1.66 \pm 0.05 = 5/3 \text{ ----- (1)}$$

Part-II: $\text{H}_2\text{O}_2/\text{MnO}_4^-$



The above an unbalanced redox chemical reaction of hydrogen peroxide with KMnO_4 is presented in which m and n are the stoichiometric coefficients of reducing and oxidising agent respectively. Here the unbalanced redox reaction may be employed as,



m x moles of reductant $\text{H}_2\text{O}_{2(l)} = n$ x moles of oxidant MnO_4^-

$$\frac{m}{n} = \frac{\text{moles of an oxidant}}{\text{moles of reductant}}$$

$$\frac{m}{n} = \frac{\text{moles of an oxidant MnO}_4^-}{\text{moles of reductant H}_2\text{O}_2}$$

$$w = N \times m \times v$$

$$\frac{m}{n} = \frac{N(\text{MnO}_4^-) \times v(\text{Volume of MnO}_4^-)}{N(\text{H}_2\text{O}_2) \times v(\text{Volume of H}_2\text{O}_2)}$$

$$\frac{m}{n} = \frac{W(\text{MnO}_4^-) \times M(\text{H}_2\text{O}_2)}{M(\text{MnO}_4^-) \times W(\text{H}_2\text{O}_2)}$$

w (Weight of a compound) = N (Normality) x m (molar mass) x v (Volume of a solution)

students performed the experiments for the above redox reaction, in which get the following (m/n) ratio here also known the Normality of Oxidant (MnO_4^-) and Reductant (H_2O_2) and Volume of H_2O_2 which was taken initially. However, at the equivalence point of the redox reaction the volume of MnO_4^- was taken. The following results was generated

$$\frac{m}{n} = 2.5 \pm 0.05 = 5/2 \text{ ----- (2)}$$

RESULT AND DISCUSSION

Two redox reactions were performed by taking same oxidant (oxidising agent) KMnO_4 , according to the stoichiometric ratio of the above two redox reaction as shown in equation (1) and

(2) it will be possible to balancing the redox reaction in acidic as well as basic medium. The results are obtained as below

Part-I




The stoichiometric ratio for the above redox reaction is $m/n = 5/3$, which is replaced in the reaction,



The question is arising how to balanced the redox reaction in acidic as well as in basic medium?

In *Acidic medium* to complete balancing the reaction in which oxygen and hydrogen must be balanced. One oxygen becomes equal to the one water molecule but water molecules must be added on opposite side of the oxygen atom, here adding 12H₂O molecules on the right-hand side. To balancing, finally adding hydrogen atoms and getting the complete balanced redox reaction.



In *Basic medium* to complete balancing the reaction in which the water molecules must be added on same side of the oxygen atom, here adding 12H₂O molecules on the left-hand side. To balancing, finally adding hydroxide ions (OH⁻) ions and getting the complete balanced redox reaction.



Part-II




The stoichiometric coefficient ratio for the redox reaction is $m/n = 5/2$, which is mentioned in the redox reaction,




Similar method which is mentioned in the part I is applying for balancing the redox reaction in acidic as well as in basic medium.

In *Acidic medium*: adding 8H₂O molecules on the right-hand side using hydrogen atoms for complete balancing redox reaction.



In *Basic medium* to complete balancing the reaction in which the water molecules must be

added on same side of the oxygen atom, here adding 12H₂O molecules on the left-hand side. To balancing, finally adding hydroxide ions (OH⁻) ions and getting the complete balanced redox reaction. The final complete balance redox reaction in basic medium is



CONCLUSION

This is the sophisticated method or procedure to balancing the redox reaction in acidic as well as in basic medium after performing the experimental redox titration method, more work is needed other than the oxidising agent (KMnO₄⁻) for the different types of redox titrimetric methods for balancing the reactions.

REFERENCES

1. Aliyar Mausavi. Revival of the oxidation number method for balancing redox equations. *Transactions of the Royal Society of South Africa*. Volume 73, 2018 - Issue 1.
2. Abraham Tamir. Conservation Law of Mass. *J Chem Eng Process Technol* 2013, 4:8.
3. Balakrishnan Viswanathan Mohamed Shajahan Gulam Razul. A Method for Teaching How to Balance Redox Reactions by Building Up Molecules. *World Journal of Chemical Education*. 2020, 8(2), 67-70.
4. B. R. Puri, L.R. Sharma, M.S. Pathania & Navjot Kaur. Electrochemistry Part-II Electromotive force of Galvanic cell. *Fundamentals of physical chemistry*. 44th Edition, pp-782 -820.
5. Eni Generalić and Nives Vladislavić. Aggregate Redox Species Method - An Improved Oxidation Number Change Method for Balancing Redox Equations. *Chemistry Journal* Vol. 4, No. 3, 2018, pp. 43-49.
6. Hans-Dieter Barke. Two ideas of the redox reaction: misconceptions and their challenge in chemistry education. *AJCE*, 2012, 2(2).
7. Matthew S. Fox. On Balancing Acidic and Basic Reduction/Oxidation Reactions with a Calculator. *World Journal of Chemical Education*, 2015, Vol. 3, No. 3, 74-77.

8. Roger L. DeKock, Benjamin M. Brandsen. Balancing Redox reaction. *J. Chem. Educ.* 2010, 87, 5, 476-477.
9. Vikram R. Jadhav. Mathematical treatment to understanding the concentration terms. *International Journal of Research & Review.* 2019; Vol.6; (1).
10. Peter Atkins, Tina Overton, Jonathan Rourke, Mark Weller and Fraser Armstrong. Oxidation and Reduction. *Shriver and Atkins Inorganic Chemistry.* 4th edition, pp 141-168.

How to cite this article: Authors' name upto 3 followed by et. al for more than 3. Full title of article is written here. *International Journal of Research and Review.* 2020; 7(1):1-10.
